

Quantum Theory of Condensed Matter I


Prof. Milena Grifoni
 Dr. Andrea Donarini
 Sebastian Pfaller

5.1.01 Mondays 10:15
 9.2.01 Tuesdays 12:15

Sheet 3

1. Band structure and Fermi surface of a square lattice

Consider a two-dimensional square crystal, with the ions at coordinates $\vec{R} = n\vec{a}_1 + m\vec{a}_2$, where $a = |\vec{a}_1| = |\vec{a}_2|$ is the lattice constant, and n and m are integers.



1. What is the Brillouin zone of the system?

(1 Point)

Let us assume one atomic orbital ν per atom. The tight-binding Hamiltonian in the $|\nu\vec{R}\rangle$ basis can be written as

$$\hat{h} = \epsilon_\nu \sum_{\vec{R}} |\nu\vec{R}\rangle \langle \nu\vec{R}| - t \sum_{\substack{\vec{R}\vec{R}' \\ \text{nearest} \\ \text{neighbours}}} |\nu\vec{R}\rangle \langle \nu\vec{R}'| - t' \sum_{\substack{\vec{R}\vec{R}' \\ \text{next nearest} \\ \text{neighbours}}} |\nu\vec{R}\rangle \langle \nu\vec{R}'|,$$

or, in other words, $\gamma(\vec{a}_i) = -t$ and $\gamma(\pm(\vec{a}_1 \pm \vec{a}_2)) = -t'$.

2. Calculate the band structure for this Hamiltonian, assuming $t' = 0$ (use the LCAO secular approximation). How many bands do you find? (2 Points)

3. Assume $t' \neq 0$. How is the band structure modified with respect to the case from 1.2? (1 Point)

4. Prove that in the case of $t' = 0$ the Fermi energy is zero for a density of (valence) electrons corresponding to one electron per atom. Calculate analytically the form of the Fermi surface for this case. (2 Points)

5. Calculate the Fermi surface for very low electron density: namely for Fermi energies $\epsilon_F = -4(t + t') + \delta$, with $\delta \ll t$. (2 Points)


6. Sketch the Fermi surface in the following 3 cases (you can use e.g. Maple, Mathematica, Matlab...)

- $t' = 0, \epsilon_F = 0$ (just a check of the analytics!)
- $t' = 0, \epsilon_F = -0.2t$
- $t' = 0.1t, \epsilon_F = -0.2t$

(2 Points)

2. Diatomic square lattice (Oral)

Consider now a crystal shown below, with two atoms in a unit cell (\bullet and \circ), and one orbital ν per atom.



1. Draw the first Brillouin zone for the system.
2. Write down the tight-binding Hamiltonian in the LCAO basis $\{|\nu_{\bullet/o}\vec{R}\rangle\}$, where $\varphi_{\nu(\bullet/o)}(\vec{r} - \vec{R} - \vec{\tau}_{\bullet/o}) := \langle \vec{r} | \nu_{\bullet/o}\vec{R} \rangle$, assuming that only overlap integrals between nearest neighbours are not zero. Calculate the band structure for this Hamiltonian (use the LCAO secular approximation). How many bands do you find?
3. Consider the system with the valence electron density corresponding to one electron per ion: is it a metal, a semiconductor, or an insulator? Explain why.
4. In the case of identical atoms, where the on-site energies are equal ($\epsilon_{\nu\bullet} = \epsilon_{\nu o}$), the model studied here reduces to the one in Problem 1.2 in the limit $t' = 0$. Prove that the band structures calculated in the two exercises coincide.
Hint: How do the different Brillouin zones map into each other?

Frohes Schaffen!